

CLENPIQ PREP

Split Dose



(501) 664-6980 | GastroArkansas.com

Gastro-Intestinal Center
10001 Lile Dr
Little Rock, AR 72205
(501) 663-1074
Fax: (501) 663-0906

Endoscopy Center of Little Rock
4200 N. Rodney Parham Rd.
Suite 203
Little Rock, AR 72212
(501) 228-4445
Fax: (501) 228-0110

GastroArkansas Endoscopy of Conway
455 Hogan Ln.
Conway, AR 72034
(501) 764-1960
Fax: (501) 664-6081

Report to the outpatient department at

Date: _____ **Time:** _____

We ask for 3 day notice to cancel your procedure. Missed appointments or cancellations without 3 day notice will incur a \$100 fee. Please bring your drivers license and insurance card to your appointment.

ONE WEEK BEFORE YOUR PROCEDURE

Do not take IRON, vitamins or herbal supplements. Continue to take your usual medications UNLESS stated by our physician. Do not take fish oil 2 days prior to procedure. **Avoid food with seeds (flax seed, popcorn, corn, nuts, berries, etc.)**

If you are taking glucagon like peptides receptor agonists such as Ozempic, Mounjaro, Semaglutide, and others in this category, please hold this medication for 7 days prior to procedure.

DAY BEFORE YOUR PROCEDURE

DO NOT EAT ANY SOLID FOODS THE DAY BEFORE YOUR TEST

Stay on clear liquids the entire day prior to procedure. It is important that you stay hydrated by using a variety of clear liquids from the list below.

- Bouillon (99% fat free) and Broth
- Apple Juice
- Plain coffee (no cream)
- Popsicles (no red or purple)
- Jello (no red or purple)
- Ginger Ale, Water, Sprite, 7-UP
- Soft drinks (Coke, Diet Coke, Dr. Pepper, etc.)
- Plain tea (no flavoring)
- Hard candy (such as sour balls)

NO MILK OR MILK PRODUCTS - Including Coffee Mate

NO PULP

NO RED OR PURPLE PRODUCTS

DIABETICS: Reduce your insulin from _____ units of _____ in the AM to _____ units. In the PM, reduce your insulin from _____ units of _____ to _____ units. If you take oral diabetic medications – do NOT take them the morning of your procedure. Please check your blood sugar before you leave home and let the nurse know the results before the procedure. Please monitor your blood sugar while prepping for the procedure.

ESSENTIAL MEDICATIONS: On the morning of the procedure, DO take your heart, blood pressure, seizure, anxiety, and depression medications with a small sip of water before 5am.

BLOOD THINNERS: If you take antiplatelet agents or anticoagulants (blood thinners) such as Plavix/ Clopidogrel, Effient/ Prasugrel, Brilinta/ Ticagrelor, Pradaxa/ Dabigatran, Xarelto/ Rivaroxaban, Eliquis/ Apixaban, Coumadin/Warfarin, etc., your physician will give you instructions on holding these medications prior to procedure. You do not need to stop aspirin.

** Hold your _____ for _____ days prior to your procedure.

*Please see reverse side

CLENPIQ PREP

Split Dose


The Clenpiq Prep Carton contains 2 bottles (5.4 oz each) and an 8 oz cup for drinking clear liquids. You must complete the entire prep to ensure the most effective cleansing.



ON THE DAY BEFORE YOUR COLONOSCOPY, stop eating all solid food and dairy, and start hydrating by drinking clear liquids.


Evening Before

At 8pm



Drink one bottle of CLENPIQ right from the bottle.
Follow with at least 40 oz (5 cups) of clear liquids.
Finish liquids over the next 5 hours.

Morning of



Drink one bottle of CLENPIQ right from the bottle.
Follow with at least 24 oz (3 cups) of clear liquids.
Finish liquids within 1 hour.

- **You must follow the instructions carefully to ensure your system will be clean, clear, and ready for examination. If instructions are not followed, the physician may be unable to perform your procedure.**
- **Nothing by mouth after midnight other than your prep, this includes gum, candy, mints, and all tobacco products.**
- **Someone must accompany you to your procedure to drive you home. You will be sedated and unable to drive for the remainder of the day.**
- **Failure to follow these instructions will result in a delay of your procedure time.**

If you have questions, please call our office at (501) 664-6980.

You will be billed separately from the Out-Patient Facility.