# MIRALAX PREP Split Dose



(501) 664-6980 | GastroArkansas.com

# **Gastro-Intestinal Center**

10001 Lile Dr Little Rock, AR 72205 (501) 663-1074 Fax: (501) 663-0906

# Endoscopy Center of Little Rock

4200 N. Rodney Parham Rd. Suite 203 Little Rock, AR 72212 (501) 228-4445 Fax: (501) 228-0110

# GastroArkansas Endoscopy of Conway

455 Hogan Ln. Conway, AR 72034 (501) 764-1960 Fax: (501) 664-6081

| Report to the outpat | ient department at   |
|----------------------|--|
| Date:                | Time:  |
| _                    | cancel your procedure. Missed appointments or lay notice will incur a \$100 fee. Please bring your drivers rd to your appointment. |

## ONE WEEK BEFORE YOUR PROCEDURE

Do not take IRON, vitamins or herbal supplements. Continue to take your usual medications UNLESS stated by our physician. Do not take fish oil 2 days prior to procedure.

Avoid food with seeds (flax seed, popcorn, corn, nuts, berries, etc.)

If you are taking glucagon like peptides receptor agonists such as Ozempic, Mounjaro, Semaglutide, and others in this category, please hold this medication for 7 days prior to procedure.

# DAY BEFORE PROCEDURE DAY BEFORE YOUR PROCEDURE

### DO NOT EAT ANY SOLID FOODS THE DAY BEFORE YOUR TEST

Stay on clear liquids the entire day prior to procedure. It is important that you stay hydrated by using a variety of clear liquids from the list below.

- Bouillon (99% fat free) and Broth
- Apple Juice
- Plain coffee (no cream)
- Popsicles (no red or purple)
- Jello (no red or purple)

- Ginger Ale, Water, Sprite, 7-UP
- Soft drinks (Coke, Diet Coke, Dr. Pepper, etc.)
- Plain tea (no flavoring)
- Hard candy (such as sour balls)

# NO MILK OR MILK PRODUCTS - Including Coffee Mate NO PULP

# **NO RED OR PURPLE PRODUCTS**

| <b>DIABETICS:</b> Reduce your insulin from  | units of         | in the AM to                     | units. In the PM, reduce         |
|---|------------------|----------------------------------|----------------------------------|
| your insulin fromunits of   | to               | units. If you take oral diabetic | medications – do <b>NOT</b> take |
| them the morning of your procedure. Pleas results before the procedure. Please monitors | •                | <u> </u>                         |                                  |
| ESSENTIAL MEDICATIONS: On the mornin  | g of the procedu | ıre, DO take your heart, blood p | oressure, seizure, anxiety, and  |

**ESSENTIAL MEDICATIONS:** On the morning of the procedure, DO take your heart, blood pressure, seizure, anxiety, and depression medications with small sip of water **before 5am**.

**BLOOD THINNERS:** If you take antiplatelet agents or anticoagulants (blood thinners) such as Plavix/ Clopidogrel, Effient/ Prasugrel, Brilinta/ Ticagrelor, Pradaxa/ Dabigatran, Xarelto/ Rivaroxaban, Eliquis/ Apixaban, Coumadin/Warfarin, etc., your physician will give you instructions on holding these medications prior to procedure. You do not need to stop aspirin.

| ** Hold your days prior to | your proced | dure |
|----------------------------|-------------|------|
|----------------------------|-------------|------|

# MIRALAX PREP Split Dose



(501) 664-6980 | GastroArkansas.com

# **DAY BEFORE PROCEDURE**

# DO NOT EAT ANY SOLID FOODS THE DAY BEFORE YOUR TEST

• You must consume 8 ounces of clear liquids every hour.

## YOU WILL NEED THE FOLLOWING ITEMS:

- 4 Dulcolax Laxative Tablets
- 238 gram bottle of MiraLax
- 64 ounce bottle of Gatorade (No Red, Orange, or Purple)
- May use (2) 32 ounce bottles for different flavors
  - 3PM: Take the 4 Dulcolax Laxative Tablets
  - 8PM: Mix the 238 gram bottle of MiraLax in 64 ounces of Gatorade. Shake the solution until MiraLax is dissolved. Drink half of the MiraLax Gatorade solution every 10-15 minutes until it is gone.

### DAY OF PROCEDURE

- At \_\_\_\_\_: Drink the other half of the MiraLax Gatorade solution, every 10 15 minutes until it is gone.
- NEXT: Drink 16 ounces of clear liquid over the next hour. Do not consume anything by mouth following this step. This includes gum, candy, mints, and all tobacco products.
- Nothing by mouth after midnight other than your prep, this includes gum, candy, mints, and all tobacco products.
- You must follow the instructions carefully to ensure your system will be clean, clear, and ready for examination. If instructions are not followed, the physician may be unable to perform your procedure.
- Someone must accompany you to your procedure to drive you home. You will be sedated and unable to drive for the remainder of the day.
- Failure to follow these instructions will result in a delay of your procedure time.

If you have questions, please call our office at (501) 664-6980.